

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>7</sup> : <b>A61K 38/00</b>		<b>A2</b>	(11) International Publication Number: <b>WO 00/06185</b>
			(43) International Publication Date: 10 February 2000 (10.02.00)
(21) International Application Number: PCT/US99/17294 (22) International Filing Date: 29 July 1999 (29.07.99) (30) Priority Data: 09/126,525      30 July 1998 (30.07.98)      US 60/094,693      30 July 1998 (30.07.98)      US (63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Applications US      60/094,693 (CON) Filed on      30 July 1998 (30.07.98) US      09/126,525 (CON) Filed on      30 July 1998 (30.07.98) (71) Applicant (for all designated States except US): BIOMEASURE INCORPORATED [US/US]; 27 Maple Street, Milford, MA 01757-3650 (US). (72) Inventor; and (75) Inventor/Applicant (for US only): MOREAU, Jacques-Pierre [US/US]; 159 Westboro Road, Upton, MA 01568 (US). (74) Agent: TSAO, Y., Rocky; Fish & Richardson P.C., 225 Franklin Street, Boston, MA 02110-2804 (US).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>	
(54) Title: METHODS OF USING A SOMATOSTATIN ANALOGUE			
(57) Abstract			
<p>The present invention is directed to a method of treating one or more of the following disease and/or conditions, which comprises administering to a patient in need thereof the compound H-g(b)-D-Nal-Cys-Tyr-D-Trp-Lys-Val-Cys-Thr-NH<sub>2</sub>, where the Cysteines are bonded by a disulfide bond, or a pharmaceutically acceptable salt thereof, most preferably the acetate salt of the compound, in the treatment of certain diseases and/or conditions such as gastroenterological conditions and/or diseases, endocrinological diseases and/or conditions, various types of cancers and conditions associated with cancer such as cancer cachexia and in the treatment of hypotension and panic attacks.</p>			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NR	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## METHODS OF USING A SOMATOSTATIN ANALOGUE

### 5. Background of the Invention

The present invention is directed to a method of treating one or more of the following diseases and/or conditions in a patient in need thereof, which comprises the administration of the compound of the formula H- $\beta$ -D-Nal-Cys-Tyr-D-Trp-Lys-Val-Cys-Thr-NH<sub>2</sub> (also known as lanreotide), where the two  
10 Cysteines are bonded by a disulfide bond, or a pharmaceutically acceptable salt thereof, most preferably the acetate salt of the compound, in the treatment of certain diseases and/or conditions such as gastroenterological conditions and/or diseases, such as Crohn's disease, systemic sclerosis, external and internal pancreatic pseudocysts and ascites, ViPoma, nesidoblastosis, hyperinsulinism,  
15 gastrinoma, Zollinger-Ellison Syndrome, diarrhea, AIDS related diarrhea, chemotherapy related diarrhea, scleroderma, Irritable Bowel Syndrome, pancreatitis, upper gastrointestinal bleeding, postprandial portal venous hypertension especially in cirrhotic patients, complications of portal hypertension, small bowel obstruction, gastroesophageal reflux, duodenogastric  
20 reflux and in treating endocrinological diseases and/or conditions, such as Cushing's Syndrome, gonadotropinoma, hyperparathyroidism, Graves' Disease, diabetic neuropathy, macular degeneration, hypercalcemia of malignancy, Paget's disease, and polycystic ovary disease; in treating various types of cancer such as thyroid cancer, leukemia, meningioma and conditions associated  
25 with cancer such as cancer cachexia; in the treatment of such conditions as hypotension such as orthostatic hypotension and postprandial hypotension and panic attacks.

Lanreotide is an analog of somatostatin and is known to inhibit growth hormone release as well as inhibit insulin, glucagon and pancreatic exocrine  
30 secretion.

U.S. Patent No. 4,853,371 discloses lanreotide, a method for making it and a method for inhibiting the secretion of growth hormone, insulin, glucagon and pancreatic exocrine secretion.

U.S. Patent No. 5,147,856 discloses the use of lanreotide of treating  
35 restenosis.

U.S. Patent No. 5,411,943 discloses the use of lanreotide for treating hepatoma.

U.S. Patent No. 5,073,541 discloses the use of lanreotide for treating lung cancer.

5 U.S. Application No. 08/089,410 filed July 9, 1993 discloses the use of lanreotide for treating melanoma.

U.S. Patent No. 5,504,069 discloses the use of lanreotide for inhibiting the accelerated growth of a solid tumor.

10 U.S. Application No. 08/854,941 filed May 13, 1997, discloses the use of lanreotide for decreasing body weight.

U.S. Application No. 08/854,943 filed May 13, 1997, discloses the use of lanreotide for treating insulin resistance and Syndrome X.

U.S. Patent No. 5,688,418 discloses the use of lanreotide for prolonging the survival of pancreatic cells.

15 PCT Application No. PCT/US97/14154 discloses the use of lanreotide for treating fibrosis.

U.S. Application No. 08/855,311 filed May 13, 1997, discloses the use of lanreotide for treating hyperlipidemia.

20 U.S. Application No. 08/440,061 filed May 12, 1995, discloses the use of lanreotide for treating hyperamylinemia.

U.S. Application No. 08/852,221 filed May 7, 1997, discloses the use of lanreotide for treating hyperprolactinemia and prolactinomas.

The contents of the foregoing patents and applications are incorporated herein by reference.

## 25 Summary of the Invention

This invention is directed to a method of treating a disease or condition which comprises administering to a patient in need thereof an effective amount of the compound H- $\beta$ -D-Nal-Cys-Tyr-D-Trp-Lys-Val-Cys-Thr-NH<sub>2</sub>, where the two Cysteines are bonded by a disulfide bond, or a pharmaceutically acceptable salt thereof, wherein the disease or condition is selected from the group consisting of  
30 systemic sclerosis, pancreatic pseudocysts, pancreatic ascites, VIPoma, nesidoblastosis, hyperinsulinism, gastrinoma, Zollinger-Ellison Syndrome, hypersecretory diarrhea, scleroderma, irritable bowel syndrome, upper gastrointestinal bleeding, postprandial portal venous hypertension, complications  
35 of portal hypertension, small bowel obstruction, duodenogastric reflux, Cushing's

Syndrome, gonadotropinoma, hyperparathyroidism, diabetic neuropathy, macular degeneration, hypercalcemia of malignancy, Paget's disease, meningioma, cancer cachexia, psoriasis, hypotension and panic attacks.

5 A preferred method of the immediately foregoing method is where the acetate salt of H- $\beta$ -D-Nal-Cys-Tyr-D-Trp-Lys-Val-Cys-Thr-NH<sub>2</sub> is administered.

A preferred method of the immediately foregoing method is where the disease or condition is selected from the group consisting of VIPoma, nesidoblastosis, hyperinsulinism, gastrinoma, hypersecretory diarrhea, irritable bowel syndrome, upper gastrointestinal bleeding, postprandial portal venous  
10 hypertension, especially in cirrhotic patients, complications of portal hypertension, small bowel obstruction, diabetic neuropathy, meningioma and cancer cachexia.

A preferred method of the immediately foregoing method is where the disease or condition treated is selected from the group consisting of VIPoma, nesidoblastosis, hypersecretory diarrhea, irritable bowel syndrome, small bowel  
15 obstruction and diabetic neuropathy.

In another aspect, the present invention is directed to a pharmaceutical composition comprising a pharmaceutically acceptable carrier and an effective amount of the acetate salt of H- $\beta$ -D-Nal-Cys-Tyr-D-Trp-Lys-Val-Cys-Thr-NH<sub>2</sub> to  
20 treat a disease or condition wherein the disease or condition is selected from the group consisting of systemic sclerosis, pancreatic pseudocysts, pancreatic ascites, VIPoma, nesidoblastosis, hyperinsulinism, gastrinoma, Zollinger-Ellison Syndrome, hypersecretory diarrhea, scleroderma, irritable bowel syndrome, upper gastrointestinal bleeding, postprandial portal venous hypertension, especially in cirrhotic patients, complications of portal hypertension, small bowel  
25 obstruction, duodenogastric reflux, Cushing's Syndrome, gonadotropinoma, hyperparathyroidism, diabetic neuropathy, macular degeneration, hypercalcemia of malignancy, Paget's disease, meningioma, cancer cachexia, psoriasis, hypotension and panic attacks.

30

#### Detailed Description

Lanreotide is readily prepared according to the procedure disclosed in U.S. Patent No. 4,853,371, or the procedure disclosed in U.S. Patent No. 5,411,943, the teachings of which are incorporated herein by reference.

Lanreotide is currently marketed as the acetate salt in a 30 mg long-acting form and is available from Ipsen Biotech, Paris, France.

As is well known to those skilled in the art, the known and potential uses of somatostatin are varied and multitudinous. Somatostatin is known to be useful in the treatment of the diseases and/or conditions listed hereinbelow. The varied uses of somatostatin may be summarized as follows: Cushing's Syndrome (see Clark, R.V. et al, Clin. Res. 38, p. 943A, 1990); gonadotropinoma (see Ambrosi B., et al., Acta Endocr. (Copenh.) 122, 569-576, 1990); hyperparathyroidism (see Miller, D., et al., Canad. Med. Ass. J., Vol. 145, pp. 227-228, 1991); Paget's disease (see, Palmieri, G.M.A., et al., J. of Bone and Mineral Research, 7, (Suppl. 1), p. S240 (Abs. 591), 1992); VIPoma (see Koberstein, B., et al., Z. Gastroenterology, 28, 295-301, 1990 and Christensen, C., Acta Chir. Scand. 155, 541-543, 1989); nesidioblastosis and hyperinsulinism (see Laron, Z., Israel J. Med. Sci., 26, No. 1, 1-2, 1990, Wilson, D.C., Irish J. Med. Sci., 158, No. 1, 31-32, 1989 and Micic, D., et al., Digestion, 16, Suppl. 1.70. Abs. 193, 1990); gastrinoma (see Bauer, F.E., et al., Europ. J. Pharmacol., 183, 55 1990); Zollinger-Ellison Syndrome (see Mozell, E., et al., Surg. Gynec. Obstet., 170, 476-484, 1990); hypersecretory diarrhea related to AIDS and other conditions (due to AIDS, see Cello, J.P., et al., Gastroenterology, 98, No. 5, Part 2, Suppl., A163 1990; due to elevated gastrin-releasing peptide, see Alhindawi, R., et al., Can. J. Surg., 33, 139-142, 1990; secondary to intestinal graft vs. host disease, see Bianco J.A., et al., Transplantation, 49, 1194-1195, 1990; diarrhea associated with chemotherapy, see Petrelli, N., et al., Proc. Amer. Soc. Clin. Oncol., Vol. 10, P 138, Abstr. No. 417 1991); irritable bowel syndrome (see O'Donnell, L.J.D., et al., Aliment. Pharmacol. Therap., Vol. 4, 177-181, 1990); pancreatitis (see Tulassay, Z., et al., Gastroenterology, 98, No. 5, Part 2, Suppl., A238, 1990); Crohn's Disease (see Fedorak, R.N., et al., Can. J. Gastroenterology, 3, No. 2, 53-57, 1989); systemic sclerosis (see Soudah, H., et al., Gastroenterology, 98, No. 5, Part 2, Suppl., A129, 1990); thyroid cancer (see Modigliani, E., et al., Ann., Endocr. (Paris), 50, 483-488, 1989); psoriasis (see Camisa, C., et al., Cleveland Clinic J. Med., 57, No. 1, 71-76, 1990); hypotension (see Hoeldtke, R.D., et al., Arch. Phys. Med. Rehabil., 69, 895-898, 1988 and Kooner, J.S., et al., Brit. J. Clin. Pharmacol., 28, 735P-736P, 1989); panic attacks (see Abelson, J.L., et al., Clin. Psychopharmacol., 10, 128-132, 1990); scleroderma (see Soudah, H., et al., Clin. Res., Vol. 39, p. 303A, 1991);

small bowel obstruction (see Nott, D.M., et al., Brit. J. Surg., Vol. 77, p. A691, 1990); gastroesophageal reflux (see Branch, M.S., et al., Gastroenterology, Vol. 100, No. 5, Part 2 Suppl., p. A425, 1991); duodenogastric reflux (see Hasler, W., et al., Gastroenterology, Vol. 100, No. 5, Part 2, Suppl., p. A448, 1991);  
5 Graves' Disease (see Chang, T.C., et al., Brit. Med. J., 304, p. 158, 1992); polycystic ovary disease (see Prelevic, G.M., et al., Metabolism Clinical and Experimental, 41, Suppl. 2, pp 76-79, 1992); upper gastrointestinal bleeding (see Jenkins, S.A., et al., Gut., 33, pp. 404-407, 1992 and Arrigoni, A., et al., American Journal of Gastroenterology, 87, p. 1311, (abs. 275), 1992);  
10 pancreatic pseudocysts and ascites (see Hartley, J.E., et al., J. Roy. Soc. Med., 85, pp. 107-108, 1992); leukemia (see Santini, et al., 78, (Suppl. 1), p. 429A (Abs. 1708), 1991); meningioma (see Koper, J.W., et al., J. Clin. Endocr. Metab., 74, pp. 543-547, 1992); and cancer cachexia (see Bartlett, D.L., et al., Surg. Forum., 42, pp. 14-16, 1991). The contents of the foregoing references  
15 are incorporated herein by reference.

Surprisingly, the Applicant has now discovered that lanreotide itself was particularly useful in treating the conditions, disorders and disease noted hereinabove.

The usefulness of lanreotide in the various disclosed new medical uses  
20 can be better understood through the results of tests relating to the treatment of upper gastrointestinal bleeding.

Lanreotide or a pharmaceutically-acceptable salt thereof can be administered by oral, parenteral (e.g., intramuscular, intraperitoneal, intravenous or subcutaneous injection, or implant), nasal, vaginal, rectal, sublingual or topical  
25 routes of administration and can be formulated with pharmaceutically acceptable carriers to provide dosage forms appropriate for each route of administration.

Solid dosage forms for oral administration include capsules, tablets, pills, powders and granules. In such solid dosage forms, the active compound is admixed with at least one inert pharmaceutically acceptable carrier such as  
30 sucrose, lactose, or starch. Such dosage forms can also comprise, as is normal practice, additional substances other than such inert diluents, e.g., lubricating agents such as magnesium stearate. In the case of capsules, tablets and pills, the dosage forms may also comprise buffering agents. Tablets and pills can additionally be prepared with enteric coatings.

Liquid dosage forms for oral administration include pharmaceutically acceptable emulsions, solutions, suspensions, syrups, the elixirs containing inert diluents commonly used in the art, such as water. Besides such inert diluents, compositions can also include adjuvants, such as wetting agents, emulsifying and suspending agents, and sweetening, flavoring and perfuming agents.

Preparations according to this invention for parenteral administration include sterile aqueous or non-aqueous solutions, suspensions, or emulsions. Examples of non-aqueous solvents or vehicles are propylene glycol, polyethylene glycol, vegetable oils, such as olive oil and corn oil, gelatin, and injectable organic esters such as ethyl oleate. Such dosage forms may also contain adjuvants such as preserving, wetting, emulsifying, and dispersing agents. They may be sterilized by, for example, filtration through a bacteria-retaining filter, by incorporating sterilizing agents into the compositions, by irradiating the compositions, or by heating the compositions. They can also be manufactured in the form of sterile solid compositions which can be dissolved in sterile water, or some other sterile injectable medium immediately before use.

Compositions for rectal or vaginal administration are preferably suppositories which may contain, in addition to the active substance, excipients such as coca butter or a suppository wax.

Compositions for nasal or sublingual administration are also prepared with standard excipients well known in the art.

The dosage of active ingredient in the compositions of this invention may be varied; however, it is necessary that the amount of the active ingredient be such that a suitable dosage form is obtained. The selected dosage depends upon the desired therapeutic effect, on the route of administration, and on the duration of the treatment. Generally, dosage levels of between 25  $\mu$ g/kg/day to 100 mg/kg/day of body weight daily are administered as a single dose or divided into multiple doses to humans and other animals, e.g., mammals, to obtain the desired therapeutic effect.

A preferred general dosage range is 250  $\mu$ g/kg/day to 5.0 mg/kg/day of body weight daily which can be administered as a single dose or divided into multiple doses.

Further, Lanreotide can be administered in a sustained release composition such as those described in the following patents. Among those formulations, 14-day or 28-day slow release formulations will be preferred. U.S.



Patent No. 5,672,659 teaches sustained release compositions comprising Lanreotide and a polyester. U.S. Patent No. 5,595,760 teaches sustained release compositions comprising Lanreotide in a gelable form. U.S. Application No. 08/929,363 filed September 9, 1997, teaches polymeric sustained release  
5 compositions comprising Lanreotide and chitosan. U.S. Application No. 08/740,778 filed November 1, 1996, teaches sustained release compositions comprising Lanreotide and cyclodextrin. U.S. Application No. 09/015,394 filed January 29, 1998, teaches absorbable sustained release compositions of Lanreotide. The contents of the foregoing patents and applications are  
10 incorporated herein by reference.

The use of immediate or of sustained release compositions depends on the type of indications aimed at. If the indication consists of an acute or over-acute disorder, a treatment with an immediate form will be preferred over the same with a prolonged release composition. On the contrary, for preventive or  
15 long-term treatments, a prolonged release composition will generally be preferred.

Typically, to the indication upper gastrointestinal bleeding will correspond an acute or over-acute treatment with a dosage of 80 to 120  $\mu$ g/day per person during approximately 5 days. After endoscopical treatment, preventive treatment  
20 against recurrence can be performed using lanreotide sustained release forms as an adjuvant to usual treatments; for this type of treatment, 14-day sustained release forms with a total dosage of approximately 30 mg lanreotide or 28-day lanreotide forms can be used.

For other indications than upper gastrointestinal bleeding, which  
25 correspond rather long term treatments, 14-day sustained release forms with a total dosage of approximately 30 mg lanreotide or 28-day lanreotide forms will be adequate.

### Claims

1. A method of treating a disease or condition which comprises administering to a patient in need thereof an effective amount of the compound H- $\beta$ -D-Nal-Cys-Tyr-D-Trp-Lys-Val-Cys-Thr-NH<sub>2</sub>, where the two Cysteines are  
5 bonded by a disulfide bond, or a pharmaceutically acceptable salt thereof, wherein the disease or condition is selected from the group consisting of systemic sclerosis, pancreatic pseudocysts, pancreatic ascites, VIPoma, nesidoblastosis, hyperinsulinism, gastrinoma, Zollinger-Ellison Syndrome, hypersecretory diarrhea, scleroderma, irritable bowel syndrome, upper  
10 gastrointestinal bleeding, postprandial portal venous hypertension, complications of portal hypertension, small bowel obstruction, duodenogastric reflux, Cushing's Syndrome, gonadotropinoma, hyperparathyroidism, diabetic neuropathy, macular degeneration, hypercalcemia of malignancy, Paget's disease, meningioma, cancer cachexia, psoriasis, hypotension and panic attacks.
- 15 2. A method according to claim 1 wherein the acetate salt of H- $\beta$ -D-Nal-Cys-Tyr-D-Trp-Lys-Val-Cys-Thr-NH<sub>2</sub> is administered.
3. A method according to claim 2 wherein the disease or condition is selected from the group consisting of VIPoma, nesidoblastosis, hyperinsulinism, gastrinoma, hypersecretory diarrhea, irritable bowel syndrome, upper  
20 gastrointestinal bleeding, postprandial portal venous hypertension, especially in cirrhotic patients, complications of portal hypertension, small bowel obstruction, diabetic neuropathy, meningioma and cancer cachexia.
4. A method according to claim 3 wherein the disease or condition treated is selected from the group consisting of VIPoma, nesidoblastosis,  
25 hypersecretory diarrhea, irritable bowel syndrome, small bowel obstruction and diabetic neuropathy.
5. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and an effective amount of the acetate salt of H- $\beta$ -D-Nal-Cys-Tyr-D-Trp-Lys-Val-Cys-Thr-NH<sub>2</sub> to treat a disease or condition wherein the  
30 disease or condition is selected from the group consisting of systemic sclerosis, pancreatic pseudocysts, pancreatic ascites, VIPoma, nesidoblastosis, hyperinsulinism, gastrinoma, Zollinger-Ellison Syndrome, hypersecretory diarrhea, scleroderma, irritable bowel syndrome, upper gastrointestinal bleeding, postprandial portal venous hypertension, especially in cirrhotic patients,  
35 complications of portal hypertension, small bowel obstruction, duodenogastric

reflux, Cushing's Syndrome, gonadotropinoma, hyperparathyroidism, diabetic neuropathy, macular degeneration, hypercalcemia of malignancy, Paget's disease, meningioma, cancer cachexia, psoriasis, hypotension and panic attacks.